

Herbaceous Ground Cover			
Site	Species	% cover class	
Kame terrace	Amphicarpea bracteata	Trace (<1%)	
	Aralia nudicaulus	Trace (<1%)	
	Aster macrophyllus	Trace (<1%)	
	Carex pennsylvanica	Trace (<1%)	
	Gaultheria procumbens	6-25%	
	Lysimachia quadrifolia	Trace (<1%)	
	Maianthemum canadense	Trace (<1%)	
	Medeola virginiana	Trace (<1%)	
	Monotropa uniflora	Trace (<1%)	
	Oryzopsis racemosa	Trace (<1%)	
	Pteridium aquilinum	1-5%	
	Smilacina racemosa	Trace (<1%)	
	Trientalis borealis	Trace (<1%)	
	Unknown spp.	Trace (<1%)	
	Unknown spp.	Trace (<1%)	
	Uvularia sessilifolia	Trace (<1%)	
	Bryant Mtn slope	Aster undulatus	Trace (<1%)
		Aster macrophyllus	Trace (<1%)
Carex pennsylvanica		1-5%	
Deschampsia flexuosa		51-75%	
Dryopteris marginalis		Trace (<1%)	
Lysimachia quadrifolia		Trace (<1%)	
Maianthemum canadense		Trace (<1%)	
Oryzopsis racemosa		Trace (<1%)	
Polygala spp.		Trace (<1%)	
Polypodium virginianum		Trace (<1%)	
Potentilla simplex		Trace (<1%)	
Solidago bicolor		Trace (<1%)	
Solidago spp.		Trace (<1%)	
Uvularia sessilifolia		6-25%	

Ridge top	Mitchella repens	1-5%
	Aquilegia canadensis	Trace (<1%)
	Aralia nudicaulus	Trace (<1%)
	Aster macrophyllus	6-25%
	Carex pennsylvanica	1-5%
	Deschampsia flexuosa	6-25%
	Dryopteris marginalis	1-5%
	Gaultheria procumbens	Trace (<1%)
	Lysimachia quadrifolia	Trace (<1%)
	Maianthemum canadense	Trace (<1%)
	Monotropa uniflora	Trace (<1%)
	Oryzopsis racemosa	Trace (<1%)
	Polygala spp.	Trace (<1%)
	Polypodium virginianum	1-5%
	Sedum telephioides	Trace (<1%)
	Solidago bicolor	Trace (<1%)
	Unknown spp.	Trace (<1%)
	Uvularia sessilifolia	Trace (<1%)

Found in all three sites

Found at the Kame Terrace and Ridge Top Sites

Found at the Bryant Mountain Slope and Ridge Top Site

Not shared between sites